

TRUNCATED QZ METHODS FOR LARGE SCALE GENERALIZED EIGENVALUE PROBLEMS*

D. C. SORENSEN†

Abstract. This paper presents three methods for the large scale generalized eigenvalue problem $\mathbf{Ax} = \mathbf{Bx}\lambda$. These methods are developed within a subspace projection framework as a truncation and modification of the QZ -algorithm for dense problems, that is suitable for computing partial generalized Schur decompositions of the pair (\mathbf{A}, \mathbf{B}) . A generalized partial reduction to condensed form is developed by analogy with the Arnoldi process. Then truncated forward and backward QZ iterations are introduced to derive generalizations of the Implicitly Restarted Arnoldi Method and the Truncated RQ method for the large scale generalized eigenvalue problem. These two methods require the accurate solution of linear systems at each step of the iteration. Relaxing these accuracy requirements forces us to introduce non-Krylov projection spaces that lead most naturally to block variants of the QZ iterations. A two-block method is developed that incorporates k approximate Newton corrections at each iteration. An important feature is the potential to utilize k matrix vector products for each access of the matrix pair (\mathbf{A}, \mathbf{B}) . Preliminary computational experience is presented to compare the three new methods.

Key words. Generalized eigenvalue problem, Krylov projection methods, Arnoldi method, Lanczos method, QZ method, block methods, preconditioning, implicit restarting.

AMS subject classifications. 65F15, 65G05.

1. Introduction. This paper presents three methods for the large scale generalized eigenvalue problem

$$(1.1) \quad \mathbf{Ax} = \mathbf{Bx}\lambda.$$

The methods are developed within a Krylov subspace projection framework as truncations of the QZ -algorithm [13] for dense problems. These techniques provide natural extensions of the Implicitly Restarted Arnoldi Method [20] and the Truncated RQ Method [21] to the generalized eigenvalue problem. Relaxing the accuracy level required for the solutions of linear systems leads naturally to a non-Krylov block projection method. This block method does not require accurate solution of shift-invert equations and makes efficient use of each matrix access by performing k matrix-vector products instead of one.

The first two methods require accurate solutions of linear systems at each step of the iteration. However, these methods are developed within a projection a framework that can accommodate inexact solves of the shift invert equations if the standard Krylov relations are relaxed. Introducing inexact solves forces us to introduce non-Krylov projection spaces. Once the Krylov property has been given up, it is natural to consider block variants of the QZ iterations. Therefore, we have developed a two-block method that incorporates k approximate Newton corrections at each iteration. An important feature is the potential to utilize k matrix vector products for each access of the matrix pair (\mathbf{A}, \mathbf{B}) .

For some time, there has been considerable interest in improving eigenvalue methods either by making better use of spectral transformation through multi-shift Rational Krylov methods [16] or by utilizing some sort of preconditioned iterative solution of these shift-invert equations at a relaxed accuracy level [11, 10, 14, 2, 19]. The ultimate goal is to achieve the enhanced convergence properties of the spectral transformation without the cost of an

*Received February 11, 1998. Accepted for publication July 22, 1998. Recommended by R. Lehoucq.

†Department of Computational and Applied Mathematics, Rice University, Houston, TX 77005-1892, (sorensen@caam.rice.edu). This work was supported in part by NSF cooperative agreement CCR-9120008, NSF contract ASC-9408795 and by ARPA contract number DAAL03-91-C-0047 (administered by the U.S. Army Research Office).

accurate direct or iterative solution of the shift-invert equations. Generalization of the Davidson method [4] to a wider class of problems has received a lot of attention and the Jacobi-Davidson method of Sleijpen and Van der Vorst [18] has emerged as an effective variant. The methods developed here (in particular the backward form of truncated QZs) have a lot in common with Ruhe's *RKS* method. This truncated backward form also appears to be closely related to the work of De Samblanx, Meerbergen and Bultheel on implicit applications of a rational filter in *RKS* [17].

The paper begins in Sections 2 and 3 with the development of simultaneous projections of the matrices \mathbf{A} and \mathbf{B} onto two subspaces to achieve a partial reduction to condensed form

$$\begin{aligned}\mathbf{A}\mathbf{V}_k &= \mathbf{W}_k\mathbf{H}_k + \mathbf{F}_k, \quad \text{with } \mathbf{W}_k^T\mathbf{F}_k = \mathbf{0}, \\ \mathbf{B}\mathbf{V}_k &= \mathbf{W}_k\mathbf{R}_k,\end{aligned}$$

through a generalized Arnoldi process. Here, \mathbf{V}_k and \mathbf{W}_k are both $n \times k$ orthogonal matrices, \mathbf{H}_k is a $k \times k$ upper Hessenberg matrix and \mathbf{R}_k is upper triangular. With this reduction, approximate generalized eigenvalues of the pair (\mathbf{A}, \mathbf{B}) are obtained from the projected pair $(\mathbf{H}_k, \mathbf{R}_k)$.

As with the standard Arnoldi process, storage and arithmetic costs are prohibitive for large k . Thus, restarting schemes are essential and two possibilities are developed. In Section 4, forward and backward variants of the implicitly shifted *QZ* iteration are developed for dense generalized problems. These are analogous to the *QR* and *RQ* iterations for the standard problem. Truncated forms of these forward and backward *QZ* -iterations are developed in Section 5. The forward form is analogous to implicit restarting [20] while the backward form generalizes the truncated *RQ* iteration [21]. These developments result in methods that are effective in computing a few (k) selected eigenvalues and corresponding eigenvectors within a fixed pre-determined storage requirement proportional to $n \cdot k$ and work proportional to $n \cdot k^2 + \mathcal{O}(k^3)$.

The generalized Arnoldi process requires the solution of a linear system at each step regardless of how it is organized. Depending on certain choices, this amounts to applying a mathematically equivalent standard Arnoldi process to one of the following matrix operators:

$$\mathbf{B}^{-1}\mathbf{A}, \quad \mathbf{A}\mathbf{B}^{-1}, \quad \text{or } (\mathbf{A} - \sigma\mathbf{B})^{-1}\mathbf{B}.$$

The backward variant of the truncated *QZ* iteration makes the most economical use of storage but tends to require more LU-factorizations than the forward variant. Very limited computational experience with all three methods shall be presented in Section 7. No reliable conclusions on comparative performance can be drawn from these limited tests.

Throughout this paper, capital and lower case Latin letters denote matrices and vectors respectively, while lower case Greek letters denote scalars. The j -th canonical basis vector is denoted by \mathbf{e}_j . The Euclidean norm is used exclusively and is denoted by $\|\cdot\|$. The transpose of a matrix \mathbf{A} is denoted by \mathbf{A}^T and the conjugate transpose by \mathbf{A}^H . Upper Hessenberg matrices will appear frequently and are usually denoted by the letter \mathbf{H} . The notation $\mathbf{M}(:, 1 : k)$ and $\mathbf{M}(1 : k, 1 : k)$ denote the leading k columns and the leading $k \times k$ principal submatrix of \mathbf{M} .

2. Subspace Projection. Certainly, projection methods are prominent for the iterative solution of linear systems and for computing a few eigenvalues of a large matrix or matrix pencil. In the case of the standard problem $\mathbf{A}\mathbf{x} = \mathbf{x}\lambda$, Krylov subspace projection results in the Lanczos/Arnoldi class of methods. These may be viewed as systematic ways to extract additional eigen-information from the sequence of vectors produced by a power iteration.

In hope to obtain additional information through various linear combinations of the power sequence, it is natural to formally consider the *Krylov* subspace

$$\mathcal{K}_k(\mathbf{A}, \mathbf{v}_1) = \text{Span} \{ \mathbf{v}_1, \mathbf{A}\mathbf{v}_1, \mathbf{A}^2\mathbf{v}_1, \dots, \mathbf{A}^{k-1}\mathbf{v}_1 \}$$

and to attempt to formulate the best possible approximations to eigenvectors from this subspace.

Approximate eigenpairs are constructed by imposing a Galerkin condition: A vector $\mathbf{x} \in \mathcal{K}_k(\mathbf{A}, \mathbf{v}_1)$ is called a *Ritz vector* with corresponding *Ritz value* θ if the Galerkin condition

$$\langle \mathbf{w}, \mathbf{A}\mathbf{x} - \mathbf{x}\theta \rangle = 0, \text{ for all } \mathbf{w} \in \mathcal{K}_k(\mathbf{A}, \mathbf{v}_1)$$

is satisfied. It is well known that the Lanczos/Arnoldi iteration computes an orthonormal basis \mathbf{V}_k for this Krylov subspace along with a small projected matrix $\mathbf{H}_k = \mathbf{V}_k^H \mathbf{A} \mathbf{V}_k$ of order k from which Ritz values and vectors may be obtained: (\mathbf{x}, θ) is a Ritz pair if and only if $\mathbf{H}_k \mathbf{y} = \mathbf{y}\theta$ and $\mathbf{x} = \mathbf{V}_k \mathbf{y}$.

Several schemes have been developed to extend the Krylov subspace idea to the generalized problem (1.1). These extensions are generally based upon a conversion of the generalized problem to a standard one. Perhaps the most successful variant [5] is to use the *spectral transformation*

$$(\mathbf{A} - \sigma \mathbf{B})^{-1} \mathbf{B} \mathbf{x} = \mathbf{x} \nu.$$

An eigenvector \mathbf{x} of this transformed problem is also an eigenvector of the original problem (1.1) with the corresponding eigenvalue given by $\lambda = \sigma + \frac{\nu}{\nu}$. In applications, \mathbf{B} is often symmetric and positive (semi-)definite and then it is helpful to work with the \mathbf{B} (semi-)inner product in the Lanczos/Arnoldi process [5, 8, 12]. With this transformation, the Lanczos/Arnoldi iteration converges very rapidly to eigenvalues near the shift σ because they are transformed to extremal well-separated eigenvalues and also because eigenvalues far from σ are damped (mapped near zero).

To utilize this transformation in a Lanczos/Arnoldi process, the repeated operation $\mathbf{w} \leftarrow \mathbf{A} \mathbf{v}$ is replaced by repeated solutions of a shift invert equation $(\mathbf{A} - \sigma \mathbf{B}) \mathbf{w} = \mathbf{B} \mathbf{v}$ at each step of the iteration. If a sparse-direct factorization of the shifted matrix $(\mathbf{A} - \sigma \mathbf{B})$ is possible, then this single factorization may be re-used at each step of the iteration. This approach is certainly the method of choice, but may not be practical or even possible in many important applications.

Although in some cases it may be effective to use a preconditioned iterative method to solve the shift-invert equations, there are a number of pitfalls to this approach. Typically, the shifted matrix is very ill-conditioned because σ is chosen to be near an eigenvalue of interest. Moreover, the shifted matrix is usually indefinite (or have indefinite symmetric part). These two conditions typically cause problems in the iterative solution of linear systems. Furthermore, these difficulties are exacerbated by the fact that each linear system must be solved to a considerably greater accuracy than the accuracy desired in the eigenvalue calculation. Otherwise, each step of the Lanczos/Arnoldi process will essentially involve a different matrix operator.

The underlying Krylov subspace projections associated with the Lanczos/Arnoldi process has a number of important approximation properties related to convergence and accuracy. Unfortunately, if it is not possible to solve the shift-invert equations accurately then these desirable properties are lost. However, it is possible to retain the projection idea in a way that generalizes the Arnoldi process when the shift invert equations can be solved accurately and yet can accommodate inaccurate solution of the shift-invert equations. To do this,

we must consider more general subspaces. The development of this projection framework is the primary topic of this paper. It is inspired by the following well known result.

LEMMA 2.1. *If \mathbf{A} and \mathbf{B} are complex matrices of order n , then there are unitary matrices \mathbf{V}, \mathbf{W} an upper Hessenberg matrix \mathbf{H} and an upper triangular matrix \mathbf{R} all of order n such that*

$$(2.1) \quad \begin{aligned} \mathbf{AV} &= \mathbf{WH}, \\ \mathbf{BV} &= \mathbf{WR}. \end{aligned}$$

This factorization can be computed in a finite number ($\mathcal{O}(n^3)$) of rational arithmetic and square root operations.

Proof. See [7]. \square

For the standard problem ($\mathbf{B} = \mathbf{I}$) this lemma reduces to the statement that \mathbf{A} may be put in condensed form by unitary similarity transformations. The Arnoldi process produces a partial reduction of \mathbf{A} to condensed (Hessenberg) form

$$\mathbf{AV}_k = \mathbf{V}_k \mathbf{H}_k + \mathbf{F}_k,$$

with $\mathbf{V}_k^T \mathbf{V}_k = \mathbf{I}_k$ and $\mathbf{V}_k^T \mathbf{F}_k = \mathbf{0}$. This may be interpreted simply as a truncation of the full reduction. It turns out that $\mathbf{F}_k = \mathbf{f}_k \mathbf{e}_k^T$ is a rank one matrix and this property is intrinsically tied to the fact that $\{\mathbf{V}_j : j = 1, 2, \dots, k\}$ is a sequence of orthonormal bases for the nested sequence of Krylov subspaces $\mathcal{K}_j(\mathbf{A}, \mathbf{v}_1)$. The Hessenberg matrix $\mathbf{H}_k = \mathbf{V}_k^T \mathbf{AV}_k$ is the orthogonal projection of \mathbf{A} onto the subspace $\mathcal{K}_k(\mathbf{A}, \mathbf{v}_1)$ as represented in the basis \mathbf{V}_k and

$$\mathbf{F}_k = (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^T) \mathbf{AV}_k.$$

If $k = n$ then $\mathbf{F}_k = \mathbf{0}$ and this provides a complete reduction of \mathbf{A} to condensed (Hessenberg) form.

The generalization suggested by Lemma (2.1) is

$$(2.2) \quad \begin{aligned} \mathbf{F}_k &= (\mathbf{I} - \mathbf{W}_k \mathbf{W}_k^T) \mathbf{AV}_k, \\ \mathbf{BV}_k &= \mathbf{W}_k \mathbf{R}_k. \end{aligned}$$

where $\mathbf{W}_k^T \mathbf{W}_k = \mathbf{V}_k^T \mathbf{V}_k = \mathbf{I}_k$. This projection makes the residual \mathbf{F}_k orthogonal to $\text{Range}(\mathbf{BW}_k)$, since the columns of \mathbf{W}_k form an orthonormal basis for that space. The Arnoldi process for the standard problem systematically produces the columns of \mathbf{V}_k , $k = 1, 2, \dots, n$ at the cost of a matrix vector product $\mathbf{y} \leftarrow \mathbf{A}\mathbf{v}$ and an orthogonal decomposition of this vector into a component in the existing Krylov space and one that is orthogonal to it.

The extension of this process to the generalized problem still requires the solution of a linear system at each step. Nevertheless, it is interesting to develop this generalized Arnoldi process, along with two restarting variants which will be developed in Sections 3,4,5. These algorithms are significant by themselves, but they may also be viewed as laying the groundwork for developing schemes that can relax the accuracy requirement on the shift-invert equations and yet retain the projection properties in the framework of a truncated reduction to condensed form.

3. Generalizing the Arnoldi reduction. The projection in equations (2.2) are well defined for any matrix \mathbf{V}_k , but it is not clear which choice of \mathbf{V}_k will provide good approximations to eigenvalues. The success of the implicitly restarted Lanczos/Arnoldi processes viewed as truncated QR iterations provides considerable motivation to develop a truncation of the QZ iteration in this projection framework.

The factorization in (2.1) provides an initial reduction of the pair (\mathbf{A}, \mathbf{B}) to an equivalent pair (\mathbf{H}, \mathbf{R}) in condensed form. This reduction precedes the *QZ* iteration just as reduction to Hessenberg form precedes the *QR* iteration. In fact, the two reductions are identical when $\mathbf{B} = \mathbf{I}$. The Arnoldi process may be derived (for $\mathbf{B} = \mathbf{I}$) simply by equating the leading k columns on both sides of (2.1). Therefore, this Arnoldi idea is easily generalized by doing the same thing when \mathbf{B} is not the identity matrix. This is fairly straightforward, but a little manipulation must be done to place this truncation within the projection framework of the previous section.

Truncating the relations (2.1) after k -steps yields

$$(3.1) \quad \begin{aligned} \mathbf{A}\mathbf{V}_k &= \mathbf{W}_k\mathbf{H}_k + \mathbf{f}_k\mathbf{e}_k^T \\ \mathbf{B}\mathbf{V}_k &= \mathbf{W}_k\mathbf{R}_k, \end{aligned}$$

with $\mathbf{V}_k, \mathbf{W}_k$ representing the leading k columns of \mathbf{V}, \mathbf{W} , $\mathbf{H}_k, \mathbf{R}_k$ representing the leading $k \times k$ principal submatrices of \mathbf{H}, \mathbf{R} and $\mathbf{f}_k = \mathbf{w}_{k+1}\gamma_{k+1,k}$ where \mathbf{w}_{k+1} is the $k+1$ -st column of \mathbf{W} and $\gamma_{k+1,k}$ is the k -th subdiagonal element of \mathbf{H} .

To advance this k -step factorization one step, the relations

$$(3.2) \quad \begin{aligned} \mathbf{A}[\mathbf{V}_k, \mathbf{v}] &= [\mathbf{W}_k, \mathbf{w}] \begin{bmatrix} \mathbf{H}_k & \mathbf{h} \\ \gamma\mathbf{e}_k^T & \alpha \end{bmatrix} + \mathbf{f}_{k+1}\mathbf{e}_{k+1}^T \\ \mathbf{B}[\mathbf{V}_k, \mathbf{v}] &= [\mathbf{W}_k, \mathbf{w}] \begin{bmatrix} \mathbf{R}_k & \mathbf{r} \\ 0 & \rho \end{bmatrix}, \end{aligned}$$

must be obtained to give the new columns $\mathbf{v}_{k+1} = \mathbf{v}$, $\mathbf{w}_{k+1} = \mathbf{w}$ and to update the matrices \mathbf{H}_{k+1} and \mathbf{R}_{k+1} .

Equating the leading k columns on both sides implies $\gamma = \|\mathbf{f}_k\|$ and $\mathbf{w} = \mathbf{f}_k/\gamma$. The direction \mathbf{v} must satisfy

$$(3.3) \quad \mathbf{B}\mathbf{v} = \mathbf{W}_k\mathbf{r} + \mathbf{w}\rho \quad \text{and} \quad \mathbf{V}_k^T\mathbf{v} = 0.$$

This implies that

$$0 = [\mathbf{V}_k^T\mathbf{B}^{-1}\mathbf{W}_k, \mathbf{V}_k^T\mathbf{B}^{-1}\mathbf{w}] \begin{bmatrix} \mathbf{r} \\ \rho \end{bmatrix}.$$

Now, $\mathbf{V}_k^T\mathbf{V}_k = \mathbf{I}_k$ and $\mathbf{B}\mathbf{V}_k = \mathbf{W}_k\mathbf{R}_k$ gives $\mathbf{V}_k^T\mathbf{B}^{-1}\mathbf{W}_k = \mathbf{R}_k^{-1}$ and thus

$$(3.4) \quad \mathbf{R}_k^{-1}\mathbf{r} = -\mathbf{V}_k^T\mathbf{B}^{-1}\mathbf{w}\rho.$$

Combining (3.2), (3.3) and (3.4) gives

$$(3.5) \quad \begin{aligned} \mathbf{v} &= \mathbf{B}^{-1}\mathbf{W}_k\mathbf{r} + \mathbf{B}^{-1}\mathbf{w}\rho \\ &= -\mathbf{V}_k\mathbf{R}_k^{-1}\mathbf{r} + \mathbf{B}^{-1}\mathbf{w}\rho \\ &= -\mathbf{V}_k\mathbf{V}_k^T\mathbf{B}^{-1}\mathbf{w}\rho + \mathbf{B}^{-1}\mathbf{w}\rho \\ &= (\mathbf{I} - \mathbf{V}_k\mathbf{V}_k^T)\mathbf{B}^{-1}\mathbf{w}\rho, \end{aligned}$$

with $\rho \equiv 1/(\|(\mathbf{I} - \mathbf{V}_k\mathbf{V}_k^T)\mathbf{B}^{-1}\mathbf{w}\|)$ so that $\mathbf{V}_k^T\mathbf{v} = 0$ and $\|\mathbf{v}\| = 1$. Now that the new \mathbf{v} has been determined, it follows that

$$\begin{bmatrix} \mathbf{h} \\ \alpha \end{bmatrix} = \begin{bmatrix} \mathbf{W}_k^T\mathbf{A}\mathbf{v} \\ \mathbf{w}^T\mathbf{A}\mathbf{v} \end{bmatrix}$$




FIG. 3.1. Generalized Arnoldi Reduction

and

$$f_{k+1} = Av - (W_k h + w\alpha).$$

This completes the update and leads to the the generalized Arnoldi process *GENARN* shown in Fig. 3.1.

Remark 1: The substitution $V_k = B^{-1}W_k R_k$ gives

$$(AB^{-1})W_k = W_k \hat{H}_k + \hat{f}_k e_k^T$$

and

$$(B^{-1}A)V_k = V_k \tilde{H}_k + \tilde{f}_k e_k^T$$

where $\hat{H}_k = H_k R_k^{-1}$, $\tilde{H}_k = R_k^{-1} H_k$ and $\hat{f}_k = f_k/\rho_{kk}$, $\tilde{f}_k = B^{-1} f_k$, are both Arnoldi processes that are mathematically equivalent to Algorithm 3.1.

Remark 2: Replacing A with B and replacing B with $A - \sigma B$ in this algorithm is mathematically equivalent to shift-invert Arnoldi method applied to $(A - \sigma B)^{-1}B$. With this substitution, the second relation in the previous remark becomes

$$(A - \sigma B)^{-1}BV_k = V_k \tilde{H}_k + \tilde{f}_k e_k^T.$$

This generalized Arnoldi iteration does nothing more than produce a partial reduction of the pair (A, B) to condensed form (H_k, R_k) . Just as with the standard Arnoldi process, there is no active mechanism to search for desired eigenvalues. However, methods that are analogous to implicit restarting [20] and truncated *RQ* [21] are possible and these shall be developed in the following section.

4. Implicitly Shifted QZ -Iterations. Forward and backward versions of implicitly shifted QZ iterations are developed here as simple extensions of the of the QR and RQ iterations. This leads naturally to truncated QZ iterations that generalize the truncated QR and RQ iterations developed in ([20, 21]).

In the following discussion, assume that there is a complete reduction of (\mathbf{A}, \mathbf{B}) to condensed form

$$\begin{aligned}\mathbf{AV} &= \mathbf{WH}, \\ \mathbf{BV} &= \mathbf{WR}.\end{aligned}$$

Forward QZ Iteration:

A forward QZ iteration may be developed from the following observations:
For a given shift μ , factor

$$(4.1) \quad \mathbf{H} - \mu\mathbf{R} = \mathbf{ZT}$$

where \mathbf{Z} is unitary and \mathbf{T} is upper triangular matrix. Now, factor

$$(4.2) \quad \mathbf{Z}^H \mathbf{R} = \mathbf{R}^+ \mathbf{Q},$$

where \mathbf{R}^+ is upper triangular and \mathbf{Q} is unitary. As with the QR iteration, it is straightforward to show that \mathbf{Z} is upper Hessenberg matrix in (4.1). Since both \mathbf{R} and \mathbf{R}^+ are upper triangular matrices, the relation (4.2) implies that \mathbf{Q}^H is also an upper Hessenberg matrix. It follows that

$$(4.3) \quad \begin{aligned}(\mathbf{A} - \mu\mathbf{B})\mathbf{V} &= \mathbf{WZT}, \\ \mathbf{BV} &= \mathbf{WZ}(\mathbf{Z}^H \mathbf{R}) = \mathbf{WZR}^+ \mathbf{Q}.\end{aligned}$$

Multiplying both sides of (4.3) on the right by \mathbf{Q}^H and rearranging terms gives

$$\begin{aligned}\mathbf{AV}^+ &= \mathbf{W}^+ \mathbf{H}^+, \\ \mathbf{BV}^+ &= \mathbf{W}^+ \mathbf{R}^+, \end{aligned}$$

where $\mathbf{V}^+ = \mathbf{VQ}^H$, $\mathbf{W}^+ = \mathbf{WZ}$ and $\mathbf{H}^+ = \mathbf{Z}^H \mathbf{H} \mathbf{Q}^H = \mathbf{TQ}^H + \mu\mathbf{R}^+$ is an upper Hessenberg matrix. This sequence of operations comprises a forward QZ step. It may be accomplished implicitly when \mathbf{Q} and \mathbf{Z} are represented as products of Givens' transformations.

From (4.3) it follows that

$$(\mathbf{A} - \mu\mathbf{B})\mathbf{v}_1 = \mathbf{WZT}\mathbf{e}_1 = \mathbf{BV}^+(\mathbf{R}^+)^{-1}\mathbf{T}\mathbf{e}_1 = \mathbf{Bv}_1^+ \tau$$

so that

$$(\mathbf{B}^{-1}\mathbf{A} - \mu\mathbf{I})\mathbf{v}_1 = \mathbf{v}_1^+ \tau$$

where τ is the (1,1) element of the upper triangular matrix $(\mathbf{R}^+)^{-1}\mathbf{T}$. Thus, the new starting vector \mathbf{v}_1^+ is the result of the application of a linear polynomial factor $(\mathbf{B}^{-1}\mathbf{A} - \mu\mathbf{I})$ to the old starting vector \mathbf{v}_1 .

Backward QZ Iteration:

A similar development leads to a backward QZ iteration:
For a given shift μ , factor

$$\mathbf{H} - \mu\mathbf{R} = \mathbf{TZ}$$




FIG. 4.1. Forward Implicitly Shifted *QZ*-iteration.

where Z is unitary and T is upper triangular. Now, factor

$$RZ^H = QR^+,$$

where R^+ is upper triangular and Q is unitary. As before, Z and Q^H are upper Hessenberg matrices. It follows that

$$\begin{aligned} (A - \mu B)VZ^H &= WT, \\ BVZ^H &= WQR^+. \end{aligned}$$

Thus

$$\begin{aligned} AV^+ &= W^+H^+, \\ BV^+ &= W^+R^+, \end{aligned}$$

where $V^+ = VZ^H, W^+ = WQ$ are unitary matrices and $H^+ \equiv Q^H H Z^H = TZ^H + \mu R^+$ is an upper Hessenberg matrix to complete the backwards *QZ* step.

This time, observe that

$$(A - \mu B)v_1^+ = WT e_1 = BVR^{-1}T e_1 = Bv_1 \tau$$

so that

$$\begin{aligned} v_1^+ &= (A - \mu B)^{-1} B v_1 \tau \\ &= (B^{-1} A - \mu I)^{-1} v_1 \tau \end{aligned}$$

where τ is the (1,1) element of $R^{-1}T$. Hence, the leading column of two successive V matrices are in an inverse iteration relationship.




FIG. 4.2. Backward Implicitly Shifted QZ -iteration.

5. Truncated Forward and Backward QZ -Iterations. With these versions of the QZ iteration, one can develop generalizations of truncated QR and RQ iterations for the generalized Arnoldi process. The truncated forward iteration will correspond to implicit restarting (truncated QR) developed in [20] while the truncated backward iteration will correspond to the truncated RQ iteration developed in [21]. These can be recovered from the methods developed here when $\mathbf{B} = \mathbf{I}$.

Assume now that there is a partial k -step reduction to condensed form

$$(5.1) \quad \begin{aligned} \mathbf{AV}_k &= \mathbf{W}_k \mathbf{H}_k + \mathbf{f}_k \mathbf{e}_k^T, \\ \mathbf{BV}_k &= \mathbf{W}_k \mathbf{R}_k, \end{aligned}$$

as in (3.1).

Truncated FQZ:

Select a shift μ and apply one forward QZ step to the projected pair $(\mathbf{H}_k, \mathbf{R}_k)$ to obtain $k \times k$ unitary upper Hessenberg matrices \mathbf{Q}_k^H and \mathbf{Z}_k and an upper triangular matrix \mathbf{T}_k such that $\mathbf{H}_k - \mu \mathbf{R}_k = \mathbf{Z}_k \mathbf{T}_k$. Completion of the FQZ step will give

$$\begin{aligned} \mathbf{H}_k \mathbf{Q}_k^H &= \mathbf{Z}_k \mathbf{H}_k^+ \\ \mathbf{R}_k \mathbf{Q}_k^H &= \mathbf{Z}_k \mathbf{R}_k^+, \end{aligned}$$

where \mathbf{H}_k^+ and \mathbf{R}_k^+ are order k upper Hessenberg and triangular matrices respectively. Then

$$(\mathbf{A} - \mu \mathbf{B}) \mathbf{V}_k = \mathbf{W}_k \mathbf{Z}_k \mathbf{T}_k + \mathbf{f}_k \mathbf{e}_k^T$$

and just as in the full iteration, equating the first column of both sides implies that

$$(\mathbf{A} - \mu \mathbf{B}) \mathbf{v}_1 = \mathbf{W}_k \mathbf{Z}_k \mathbf{T}_k \mathbf{e}_1 = \mathbf{B} \mathbf{V}_k^+ (\mathbf{R}_k^+)^{-1} \mathbf{T}_k \mathbf{e}_1 = \mathbf{B} \mathbf{v}_1^+ \tau.$$

Thus,

$$(\mathbf{B}^{-1} \mathbf{A} - \mu \mathbf{I}) \mathbf{v}_1 = \mathbf{v}_1^+ \tau,$$

where τ is the (1,1) element of $(\mathbf{R}_k^+)^{-1}\mathbf{T}_k$. Now,

$$(5.2) \quad \begin{aligned} \mathbf{A}\mathbf{V}_k^+ &= \mathbf{W}_k^+\mathbf{H}_k^+ + \mathbf{f}_k\mathbf{e}_k^T\mathbf{Q}_k^H \\ \mathbf{B}\mathbf{V}_k^+ &= \mathbf{W}_k^+\mathbf{R}_k^+ \end{aligned}$$

and since \mathbf{Q}_k^H is upper an Hessenberg matrix, it follows that the last row of \mathbf{Q}_k^H has the form $\mathbf{e}_k^T\mathbf{Q}_k^H = [\sigma\mathbf{e}_{k-1}^T, \gamma]$. Hence, the leading $k-1$ columns on both sides of (5.3) remain in a generalized Arnoldi relation

$$\begin{aligned} \mathbf{A}\mathbf{V}_{k-1}^+ &= \mathbf{W}_{k-1}^+\mathbf{H}_{k-1}^+ + \hat{\mathbf{f}}_{k-1}\mathbf{e}_{k-1}^T \\ \mathbf{B}\mathbf{V}_{k-1}^+ &= \mathbf{W}_{k-1}^+\mathbf{R}_{k-1}^+ \end{aligned}$$

where $\hat{\mathbf{f}}_{k-1} = \mathbf{W}_k^+\mathbf{e}_k\beta + \mathbf{f}_k\sigma$. Now, one additional generalized Arnoldi step may be performed to return this to an implicitly restarted k -step reduction.

Just as with the IRA iteration, this idea may be cast in the form of repeating the following steps: (1) Extend to a $k+p$ step factorization, (2) Apply p shifts with FQZ sweeps, (3) Truncate the last p -columns to return to a k step factorization. This will define a generalized implicitly restarted Arnoldi method.

Truncated BQZ:

To truncate the backwards QZ iteration, it will be necessary to derive relationships existing in column $k+1$ on both sides of (2.2). The required theory for the standard problem has been derived in [21] and this will generalize in a straightforward way to obtain a corresponding truncated backwards QZ equation. However, the details for completing a backward QZ sweep once this equation has been solved are a bit more intricate than in the TRQ iteration.

Following the development of the TRQ iteration, given a shift μ and the partial k -step reduction, the truncated BQZ is initiated by constructing vectors \mathbf{v} and \mathbf{w} of unit length that are orthogonal to the columns of \mathbf{V}_k and \mathbf{W}_k respectively, with $(\mathbf{A} - \mu\mathbf{B})\mathbf{v} \in \text{Range}([\mathbf{W}_k, \mathbf{w}])$. Then, a relation of the form

$$(5.3) \quad (\mathbf{A} - \mu\mathbf{B})[\mathbf{V}_k, \mathbf{v}] = [\mathbf{W}_k, \mathbf{w}] \begin{bmatrix} \mathbf{H}_k - \mu\mathbf{R}_k & \mathbf{h} \\ \beta\mathbf{e}_k^T & \alpha \end{bmatrix}.$$

is obtained to initiate a truncated BQZ step. To develop this further, assume for the moment that \mathbf{v} , \mathbf{w} , \mathbf{h} and α have been constructed to satisfy these relations. Let us postpone the construction of these quantities and first show how to complete the truncated BQZ step assuming that $(\mathbf{A} - \mu\mathbf{B})\mathbf{v} = \mathbf{W}_k\mathbf{h} + \mathbf{w}\alpha$. At this point, it is important to realize that the bordered Hessenberg matrix in (5.3) is precisely the leading principal submatrix that would appear if the full matrix $\mathbf{H} - \mu\mathbf{R}$ were partially factored into an RQ factorization from right to left using Givens' transformations up to the $k+1$ st column. The subsequent computations amount to arranging the remaining relations in the \mathbf{W} and \mathbf{R} matrices that would be in place had the first $n-k$ steps of a BQZ sweep been done. The idea is to anticipate this configuration and then complete the sweep in the leading k columns without ever computing the remaining $n-k$ columns of the BQZ relations.

At this point, the relationships for \mathbf{B} must be brought up to date. Equations must be derived that will keep \mathbf{B} in a triangular relation with the two basis sets. We first construct a vector \mathbf{w}^+ such that

$$\mathbf{B}\mathbf{v} = \mathbf{W}_k\mathbf{r} + \mathbf{w}^+\rho \quad \text{with} \quad \mathbf{W}_k^H\mathbf{w}^+ = 0$$

using classical Gram-Schmidt with the orthogonality correction scheme proposed in [3] Once this is done, we have

$$(5.4) \quad \mathbf{B}[\mathbf{V}_k, \mathbf{v}] = [\mathbf{W}_k, \mathbf{w}^+] \begin{bmatrix} \mathbf{R}_k & \mathbf{r} \\ 0 & \rho \end{bmatrix}.$$

From Equations (5.3) and (5.4) we may derive

$$\begin{aligned} \mathbf{A}[\mathbf{V}_k, \mathbf{v}] &= [\mathbf{W}_k, \mathbf{w}] \begin{bmatrix} \mathbf{H}_k - \mu\mathbf{R}_k & \mathbf{h} \\ \beta\mathbf{e}_k^T & \alpha \end{bmatrix} + [\mathbf{W}_k, \mathbf{w}^+] \begin{bmatrix} \mu\mathbf{R}_k & \mathbf{r}\mu \\ 0 & \rho\mu \end{bmatrix} \\ &= [\mathbf{W}_k, \mathbf{w}^+] \begin{bmatrix} \mathbf{H}_k & \mathbf{h} + \mathbf{r}\mu \\ \beta\theta\mathbf{e}_k^T & \alpha\theta + \rho\mu \end{bmatrix} + \mathbf{z}[\beta\mathbf{e}_k^T, \alpha], \end{aligned}$$

where \mathbf{w} has been written as $\mathbf{w} = \mathbf{w}^+ \theta + \mathbf{z}$ with $\mathbf{z}^H \mathbf{w}^+ = 0$.

At this point, in the full factorization, the leading principal $(k+1) \times (k+1)$ submatrices of the $\mathbf{H} - \mu\mathbf{R}$ and \mathbf{R} matrices are of the form

$$(5.5) \quad \hat{\mathbf{H}}_{k+1} - \mu\hat{\mathbf{R}}_{k+1} = \begin{bmatrix} \mathbf{H}_k - \mu\mathbf{R}_k & \mathbf{h} \\ \beta\theta\mathbf{e}_k^T & \alpha\theta \end{bmatrix}$$

and

$$\hat{\mathbf{R}}_{k+1} = \begin{bmatrix} \mathbf{R}_k & \mathbf{r} \\ 0 & \rho \end{bmatrix}.$$

To complete the *BQZ* step, factor

$$(5.6) \quad \hat{\mathbf{H}}_{k+1} - \mu\hat{\mathbf{R}}_{k+1} = \mathbf{T}_{k+1} \mathbf{Z}_{k+1}$$

where \mathbf{T}_{k+1} is an upper triangular and \mathbf{Z}_{k+1} is a unitary matrix. Now, factor

$$\mathbf{Q}_{k+1} \mathbf{R}_{k+1}^+ = \hat{\mathbf{R}}_{k+1} \mathbf{Z}_{k+1}^H,$$

where \mathbf{Q}_{k+1} is a unitary and \mathbf{R}_{k+1}^+ is an upper triangular matrix. As before, \mathbf{Z}_{k+1} and \mathbf{Q}_{k+1}^H are both upper Hessenberg matrices.

From Equations (5.5) and (5.6), we observe that $(\beta\mathbf{e}_k^T, \alpha)\mathbf{Z}_{k+1}^H = (0, \tilde{\alpha})$ where $\tilde{\alpha}\theta$ is the $(k+1, k+1)$ element of \mathbf{T}_{k+1} .

It follows that

$$\begin{aligned} (\mathbf{A} - \mu\mathbf{B})[\mathbf{V}_k, \mathbf{v}]\mathbf{Z}_{k+1}^H &= [\mathbf{W}_k, \mathbf{w}^+]\mathbf{T}_{k+1} + \mathbf{z}(0, \tilde{\alpha}) \\ \mathbf{B}[\mathbf{V}_k, \mathbf{v}]\mathbf{Z}_{k+1}^H &= [\mathbf{W}_k, \mathbf{w}^+]\mathbf{Q}_{k+1}\mathbf{R}_{k+1}^+, \end{aligned}$$

and then

$$(5.7) \quad \begin{aligned} (\mathbf{A} - \mu\mathbf{B})[\mathbf{V}_k, \mathbf{v}]\mathbf{Z}_{k+1}^H &= [\mathbf{W}_k, \mathbf{w}^+]\mathbf{Q}_{k+1}\mathbf{Q}_{k+1}^H\mathbf{T}_{k+1} + \mathbf{z}(0, \tilde{\alpha}) \\ \mathbf{B}[\mathbf{V}_k, \mathbf{v}]\mathbf{Z}_{k+1}^H &= [\mathbf{W}_k, \mathbf{w}^+]\mathbf{Q}_{k+1}\mathbf{R}_{k+1}^+. \end{aligned}$$

As in the full case, the relations

$$\mathbf{Q}_{k+1}^H \mathbf{T}_{k+1} + \mu\mathbf{R}_{k+1}^+ = \mathbf{Q}_{k+1}^H \hat{\mathbf{H}}_{k+1} \mathbf{Z}_{k+1}^H$$

hold and imply that $\mathbf{H}_{k+1}^+ \equiv \mathbf{Q}_{k+1}^H \hat{\mathbf{H}}_{k+1} \mathbf{Z}_{k+1}^H$ is an upper Hessenberg matrix. Therefore, deleting the $k+1$ -st column on both sides of (5.7) will give

$$\begin{aligned} \mathbf{A}\mathbf{V}_k^+ &= \mathbf{W}_k^+ \mathbf{H}_k^+ + \mathbf{f}_k^+ \mathbf{e}_k^T, \\ \mathbf{B}\mathbf{V}_k^+ &= \mathbf{W}_k^+ \mathbf{R}_k^+, \end{aligned}$$

where \mathbf{V}_k^+ is the matrix consisting of the leading k columns of $[\mathbf{V}_k, \mathbf{v}]\mathbf{Z}_{k+1}^H$ and \mathbf{W}_k^+ is the matrix consisting of the leading k columns of $[\mathbf{W}_k, \mathbf{w}^+]\mathbf{Q}_{k+1}$. The matrices \mathbf{R}_k^+ and \mathbf{H}_k^+ are the leading principal order k submatrices of \mathbf{R}_{k+1}^+ and \mathbf{H}_{k+1}^+ , and \mathbf{f}_k^+ is the last column of $[\mathbf{W}_k, \mathbf{w}^+]\mathbf{Q}_{k+1}$ scaled by the $(k+1, k)$ element of \mathbf{H}_{k+1}^+ .

This time, observe that Equation (5.7) implies that

$$(\mathbf{A} - \mu\mathbf{B})\mathbf{v}_1^+ = \mathbf{W}_k \mathbf{T}_k \mathbf{e}_1 = \mathbf{B} \mathbf{V}_k \mathbf{R}_k^{-1} \mathbf{T}_k \mathbf{e}_1$$

so that

$$\begin{aligned} \mathbf{v}_1^+ &= (\mathbf{A} - \mu\mathbf{B})^{-1} \mathbf{B} \mathbf{v}_1 \tau \\ &= (\mathbf{B}^{-1} \mathbf{A} - \mu\mathbf{I})^{-1} \mathbf{v}_1 \tau \end{aligned}$$

where τ is the $(1,1)$ element of $\mathbf{R}_k^{-1} \mathbf{T}_k$. Hence, just as in the full case, the leading columns of two successive \mathbf{V} matrices are in an inverse iteration relationship.

Now that the truncated BQZ step is understood, it is time to develop the truncated BQZ equation needed to construct \mathbf{v} , \mathbf{h} and α in equation (5.3), so that

$$(\mathbf{A} - \mu\mathbf{B})\mathbf{v} = \mathbf{W}_k \mathbf{h} + \mathbf{w} \alpha$$

with $\mathbf{w} = \mathbf{f}_k / \|\mathbf{f}_k\|$, $\mathbf{v}^H \mathbf{V}_k = 0$ and $\|\mathbf{v}\| = 1$. Existence and uniqueness for the case $\mathbf{B} = \mathbf{I}$ was developed in [21] and easily generalizes to this setting. Of the various possibilities developed there, the following seems most appropriated in this setting:

First, compute a solution $\hat{\mathbf{v}}$ to the equation

$$(5.8) \quad (\mathbf{A} - \mu\mathbf{B})\hat{\mathbf{v}} = \mathbf{W}_k \mathbf{t} + \mathbf{f}_k \eta$$

where $(\mathbf{t}^H, \eta)^H$ is an arbitrary $k+1$ vector. Then set

$$(5.9) \quad \mathbf{v} = (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}} \tau$$

where $\tau = 1 / \|(\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}}\|$. Now put

$$(5.10) \quad \mathbf{h} = \mathbf{W}_k^H (\mathbf{A} - \mu\mathbf{B})\mathbf{v} \quad \text{and} \quad \alpha = \mathbf{w}^H (\mathbf{A} - \mu\mathbf{B})\mathbf{v}.$$

The following lemma indicates why this will work.

LEMMA 5.1. *Assume $\mathbf{A} - \mu\mathbf{B}$ is nonsingular and that there is a partial reduction of (\mathbf{A}, \mathbf{B}) to condensed form as in (5.1). If $\mathbf{H}_k - \mu\mathbf{R}_k$ is nonsingular, put*

$$\mathbf{t} = (\mathbf{H}_k - \mu\mathbf{R}_k) \mathbf{s}$$

and choose $\eta \neq \mathbf{e}_k^T \mathbf{s}$, where \mathbf{s} is any k -vector. Otherwise, let $\mathbf{t} \neq 0$ be a left null vector so that

$$0 = \mathbf{t}^H (\mathbf{H}_k - \mu\mathbf{R}_k)$$

and choose η to be arbitrary. Let $\hat{\mathbf{v}}$ be the unique solution to (5.8). Then $0 \neq (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}}$, so the vector \mathbf{v} can be constructed by projection and normalized as in (5.9). Moreover,

$$(\mathbf{A} - \mu\mathbf{B})\mathbf{v} = \mathbf{W}_k \mathbf{h} + \mathbf{w} \alpha,$$

i.e. $(\mathbf{A} - \mu\mathbf{B})\mathbf{v} \in \text{Range}([\mathbf{W}_k, \mathbf{w}])$.

Proof. Suppose \mathbf{t} , η , and $\hat{\mathbf{v}}$ are constructed as prescribed in the hypothesis. If $0 = (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}}$, then $\hat{\mathbf{v}} = \mathbf{V}_k \mathbf{y}$ must hold for some nonzero k -vector \mathbf{y} . Now, this would imply

$$\begin{aligned} (\mathbf{A} - \mu \mathbf{B}) \hat{\mathbf{v}} &= (\mathbf{A} - \mu \mathbf{B}) \mathbf{V}_k \mathbf{y} \\ &= \mathbf{W}_k (\mathbf{H}_k - \mu \mathbf{R}_k) \mathbf{y} + \mathbf{f}_k \mathbf{e}_k^T \mathbf{y}. \end{aligned}$$

Substituting this on the left side of (5.8) and using orthogonality gives

$$(5.11) \quad (\mathbf{H}_k - \mu \mathbf{R}_k) \mathbf{y} = \mathbf{t} \quad \text{and} \quad \mathbf{e}_k^T \mathbf{y} = \eta.$$

If $\mathbf{H}_k - \mu \mathbf{R}_k$ is nonsingular, then $\mathbf{y} = \mathbf{s}$ and (5.11) would contradict the choice of η . Otherwise, the choice of \mathbf{t} as a null vector would lead to the following contradiction:

$$0 = \mathbf{t}^H (\mathbf{H}_k - \mu \mathbf{R}_k) \mathbf{y} = \mathbf{t}^H \mathbf{t} \neq 0.$$

This shows $0 \neq (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}}$, so that \mathbf{v} can be constructed by projection and normalized as in (5.9). It remains to show $(\mathbf{A} - \mu \mathbf{B}) \mathbf{v} \in \text{Range}([\mathbf{W}_k, \mathbf{w}])$. However, this follows easily from the relations

$$(5.12) \quad \begin{aligned} (\mathbf{A} - \mu \mathbf{B}) \mathbf{v} &= (\mathbf{A} - \mu \mathbf{B}) \hat{\mathbf{v}} - (\mathbf{A} - \mu \mathbf{B}) \mathbf{V}_k \mathbf{V}_k^H \hat{\mathbf{v}} \\ &= \mathbf{W}_k \mathbf{t} + \mathbf{f}_k \eta - [\mathbf{W}_k (\mathbf{H}_k - \mu \mathbf{R}_k) + \mathbf{f}_k \mathbf{e}_k^T] \mathbf{V}_k^H \hat{\mathbf{v}}. \end{aligned}$$

This completes the proof. \square

Since $(\mathbf{A} - \mu \mathbf{B})$ is nonsingular and $[\mathbf{W}_k, \mathbf{w}]$ is a unitary matrix, \mathbf{v}, \mathbf{h} and α are uniquely determined once \mathbf{t} and η have been specified. This justifies using (5.8) and (5.9) to compute them. However, it is remarkable that \mathbf{v}, \mathbf{h} and α are unique, regardless of the choice of \mathbf{t} and η as long as $0 \neq (\mathbf{I} - \mathbf{V}_k \mathbf{V}_k^H) \hat{\mathbf{v}}$. This result is a fairly straightforward modification of the results in Section 2 of [21].

Typically, $\mathbf{t} = \mathbf{e}_k$ is chosen because this corresponds to the standard Arnoldi process for $\mathbf{B} = \mathbf{I}$, but many other interesting choices are possible.

Remark: We may choose to cast (5.8) in the form

$$(5.13) \quad (\mathbf{I} - \mathbf{X} \mathbf{X}^H) (\mathbf{A} - \mu \mathbf{B}) (\mathbf{I} - \mathbf{Z} \mathbf{Z}^H) \hat{\mathbf{v}} = \mathbf{W}_k \mathbf{t} + \mathbf{f}_k \eta,$$

where $\mathbf{X} \equiv \mathbf{W}_k \mathbf{Y}$ and $\mathbf{Z} \equiv \mathbf{V}_k \mathbf{S}$ with $\mathbf{Y}^H \mathbf{Y} = \mathbf{S}^H \mathbf{S} = \mathbf{I}_j$. Here \mathbf{Y} and \mathbf{S} may be of dimension $k \times j$ for any $j = 1, 2, \dots, k$. Once $\hat{\mathbf{v}}$ is determined, (5.13) may be rearranged to obtain a relation of the form

$$(\mathbf{A} - \mu \mathbf{B}) \hat{\mathbf{v}} = \mathbf{W}_k \hat{\mathbf{t}} + \mathbf{f}_k \hat{\eta},$$

since

$$(\mathbf{X} \mathbf{X}^H) (\mathbf{A} - \mu \mathbf{B}) (\mathbf{I} - \mathbf{Z} \mathbf{Z}^H) \hat{\mathbf{v}} \in \text{Range}(\mathbf{W}_k) \quad \text{and} \quad (\mathbf{A} - \mu \mathbf{B}) (\mathbf{Z} \mathbf{Z}^H) \hat{\mathbf{v}} \in \text{Range}([\mathbf{W}_k, \mathbf{w}]).$$

Observe that there is no need to actually compute $\hat{\mathbf{t}}$ and $\hat{\eta}$. One may simply project and normalize as in (5.9) to get \mathbf{v} and then obtain \mathbf{h} and α as in (5.10).

This remark may have computational significance in case we choose to compute $\hat{\mathbf{v}}$ with an iterative method. In particular, if μ is a nearly converged Ritz value, then it may be a good idea to take $\mathbf{X} = \mathbf{W}_k \mathbf{y}$ where $\mathbf{y}^H (\mathbf{H}_k - \mu \mathbf{R}_k) = 0$, and $\mathbf{Z} = \mathbf{V}_k \mathbf{s}$ where $(\mathbf{H}_k - \mu \mathbf{R}_k) \mathbf{s} = 0$. This choice would tend to project out the near singularity of $(\mathbf{A} - \mu \mathbf{B})$ as suggested in [18] along the directions of the converging eigenvectors. Another possibility is to take $\mathbf{X} = \mathbf{W}_k$ and $\mathbf{Z} = \mathbf{V}_k$ as suggested in [21] to project out all of the current subspace. The latter choice is computationally more expensive (per iteration in the linear solve) but may have other advantages in the presence of clustered eigenvalues.




FIG. 5.1. Truncated Backward *QZ*-iteration

6. Inexact Arnoldi Processes. In the previous two sections, algorithms have been developed to generalize the Arnoldi process and to derive truncated forms of the forward and backward *QZ* iterations. Unfortunately, these algorithms require the accurate solution of linear systems. However, the accuracy requirement for computing the direction v through Steps (3.3)-(3.4) may be relaxed. A projection algorithm is still obtained but the Krylov property will be lost.

To relax the exact solution requirement indicated at Step(3.3), simply replace the computation of y from $By = w$ with $y = \text{itsol}(B, M, w)$ where M represents a preconditioner for B and *itsol* represents a few steps of a preconditioned iterative method for the solution of the linear system $By = w$. Formally, there is no accuracy requirement here and as little as one step of the iterative method may be specified. However, the rank-one nature of the residual F_k will be lost along with the Hessenberg form for H_k when this accuracy is relaxed.

Of course, there are algorithmic consequences of relaxing the accuracy requirements.




FIG. 6.1. An Inexact Arnoldi Process.

The relations (3.5) are no longer valid. Therefore, the relationship $\mathbf{B}\mathbf{v} = \mathbf{W}_k\mathbf{r} + \mathbf{w}\rho$ must be forced explicitly once the direction \mathbf{v} has been determined. The resulting algorithm *INXARN* is described in Fig. 6.1.

Generating Directions and the Newton Step:

Once the decision has been made to relax the Krylov property, a more general point of view may be taken. The sequence of vectors $\{\mathbf{v}_j\}$ may just as well be generated by some arbitrary process unrelated to the projections. Certainly, some relation to the shift-invert equations is desirable and the remainder of this discussion will focus on properties of the generated sequence $\{\mathbf{v}_j\}$ required for rapid convergence. With this end in mind, let us consider an arbitrary sequence of generated vectors $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_j, \dots\}$ and assume that these vectors are orthonormal in some convenient inner product.

Given this sequence, it is straightforward to obtain a derived sequence of orthogonal vectors $\{\mathbf{w}_j\}$ along with a sequence of projections that provide a partial reduction of the pair (\mathbf{A}, \mathbf{B}) to condensed form at each step:

$$\begin{aligned}
 \mathbf{V}_j &\leftarrow [\mathbf{V}_{j-1}, \mathbf{v}_j]; \\
 \mathbf{B}\mathbf{V}_j &= \mathbf{W}_j\mathbf{R}_j; \\
 \mathbf{A}\mathbf{V}_j &= \mathbf{W}_j\mathbf{H}_j + \mathbf{F}_j;
 \end{aligned}$$

with $\mathbf{W}_j^T\mathbf{W}_j = \mathbf{V}_j^T\mathbf{V}_j = \mathbf{I}_j$, $\mathbf{W}_j^T\mathbf{F}_j = \mathbf{0}$ as before through classical Gram Schmidt orthogonalization.

How should the sequence $\{\mathbf{v}_j\}$ be generated to achieve or to accelerate convergence of the Ritz values (eigenvalues of $(\mathbf{H}_j, \mathbf{R}_j)$) to selected eigenvalues of the pair (\mathbf{A}, \mathbf{B}) ? Certainly, it would be helpful to develop a connection with Newton's method and then perhaps modify those choices to reduce computational cost while retaining reasonable convergence properties. To this end, suppose $\mathbf{H}\mathbf{y} = \mathbf{R}\mathbf{y}\theta$ and $\mathbf{x} = \mathbf{V}\mathbf{y}$ with $\|\mathbf{x}\| = \|\mathbf{y}\| = 1$. Let $\lambda \in \sigma(\mathbf{A}, \mathbf{B})$ be the closest eigenvalue to θ and let \mathbf{q} be the corresponding eigenvector normalized so that $\mathbf{x}^H \mathbf{q} = 1$ (hence $\|\mathbf{q}\| \geq 1$).

With these assumptions, let us represent

$$\mathbf{q} = \mathbf{x} + \mathbf{z}, \quad \lambda = \theta + \delta,$$

with $\mathbf{x}^H \mathbf{z} = 0$ and derive the standard second order approximation from the relation $\mathbf{A}\mathbf{q} = \mathbf{B}\mathbf{q}\lambda$. Substituting, combining and rearranging terms gives

$$(6.1) \quad (\mathbf{A} - \theta\mathbf{B})\mathbf{z} = -(\mathbf{A} - \theta\mathbf{B})\mathbf{x} + \mathbf{B}\mathbf{x}\delta + \mathbf{B}\mathbf{z}\delta$$

At this point, several alternatives are available to approximate the correction vector \mathbf{z} . Two possibilities shall be examined here. The first of these gives the correction developed in [18, 6]. Since $\mathbf{x} = \mathbf{V}\mathbf{y}$, it follows that

$$\begin{aligned} -(\mathbf{A} - \theta\mathbf{B})\mathbf{x} + \mathbf{B}\mathbf{x}\delta &= -\mathbf{W}(\mathbf{H} - \theta\mathbf{R})\mathbf{y} - \mathbf{F}\mathbf{y} + \mathbf{W}\mathbf{R}\mathbf{y}\delta \\ &= -\mathbf{F}\mathbf{y} + \mathbf{W}\mathbf{R}\mathbf{y}\delta. \end{aligned}$$

Now, if both sides of equation (6.1) are multiplied on the left by $\mathbf{I} - \mathbf{W}\mathbf{W}^H$ the resulting equation is

$$(6.2) \quad (\mathbf{I} - \mathbf{W}\mathbf{W}^H)(\mathbf{A} - \theta\mathbf{B})(\mathbf{I} - \mathbf{x}\mathbf{x}^H)\mathbf{z} = -\mathbf{F}\mathbf{y} + (\mathbf{I} - \mathbf{W}\mathbf{W}^H)\mathbf{B}\mathbf{z}\delta,$$

since $0 = \mathbf{W}^H \mathbf{F}$ and $0 = \mathbf{x}^H \mathbf{z}$. From this, it also follows that equation (6.2) is consistent and there is a unique minimum norm solution \mathbf{z} . Hence the direction \mathbf{v} obtained by finding the minimum norm solution to

$$(\mathbf{I} - \mathbf{W}\mathbf{W}^H)(\mathbf{A} - \theta\mathbf{B})(\mathbf{I} - \mathbf{x}\mathbf{x}^H)\mathbf{v} = -\mathbf{F}\mathbf{y}$$

will assure that the second order correction is a member of the updated spaces $S_{\mathbf{V}} \equiv \text{Range}(\mathbf{V})$ and $S_{\mathbf{W}} \equiv \text{Range}(\mathbf{W})$ when \mathbf{v} is adjoined and the corresponding \mathbf{w} is obtained.

An alternative to the solution just developed is to treat equation (6.1) in a straightforward way assuming that the matrix $\mathbf{A} - \theta\mathbf{B}$ is nonsingular. Then

$$(6.3) \quad \mathbf{z} = -\mathbf{x} + (\mathbf{A} - \theta\mathbf{B})^{-1} \mathbf{B}\mathbf{x}\delta + (\mathbf{A} - \theta\mathbf{B})^{-1} \mathbf{B}\mathbf{z}\delta.$$

Now, using the facts $0 = \mathbf{x}^H \mathbf{z}$ and $0 = (\mathbf{I} - \mathbf{x}\mathbf{x}^H)\mathbf{x}$ gives

$$\mathbf{z} = (\mathbf{I} - \mathbf{x}\mathbf{x}^H)(\mathbf{A} - \theta\mathbf{B})^{-1} \mathbf{B}\mathbf{x}\delta + (\mathbf{I} - \mathbf{x}\mathbf{x}^H)(\mathbf{A} - \theta\mathbf{B})^{-1} \mathbf{B}\mathbf{z}\delta,$$

when both sides of equation (6.3) are multiplied on the left by the projection $(\mathbf{I} - \mathbf{x}\mathbf{x}^H)$. Now, the second order correction will be included in the updated spaces in the new direction \mathbf{v} obtained by finding the solution $\hat{\mathbf{z}}$ to

$$(\mathbf{A} - \theta\mathbf{B})\hat{\mathbf{z}} = \mathbf{B}\mathbf{x},$$

and then projecting and normalizing to get

$$\mathbf{z} = (\mathbf{I} - \mathbf{x}\mathbf{x}^H)\hat{\mathbf{z}} \quad \text{and} \quad \mathbf{v} = \mathbf{z}/\|\mathbf{z}\|.$$

Note the advantage here of adjoining the direction \mathbf{z} to the existing space. We do not need to explicitly compute δ in (6.3) as would be needed in an explicit Newton method. This projection process assures that the Newton correction is in the updated subspace so that the new Ritz vector and Ritz value will be at least as good as those obtained through an explicit Newton step.

The methods of Davidson [4], Olsen et. al, [15], Sleijpen and Van der Vorst [18] and those introduced and discussed by Knyazev [10] can all be placed within this Newton-like framework.

Blocked Formulation: Further consideration of the previous development would suggest that a block formulation is more appropriate than a single vector approach when the Krylov property is no longer enforced. To develop this, we assume a partial decomposition of the form

$$(6.4) \quad \begin{aligned} \mathbf{A}\mathbf{V}_1 &= \mathbf{W}_1\mathbf{H}_{11} + \mathbf{F}_1, \quad \text{with } \mathbf{W}_1^H\mathbf{F}_1 = 0, \\ \mathbf{B}\mathbf{V}_1 &= \mathbf{W}_1\mathbf{R}_{11}, \end{aligned}$$

where $\mathbf{V}_1, \mathbf{W}_1, \mathbf{F}_1$ are $n \times k$ matrices and $\mathbf{H}_{11}, \mathbf{R}_{11}$ are $k \times k$ matrices. We then construct the $n \times k$ matrix \mathbf{V}_2 as follows:

$$\begin{aligned} \mathbf{V} &= (\mathbf{I} - \mathbf{V}_1\mathbf{V}_1^H)p(\mathbf{A}, \mathbf{B})\mathbf{F}_1, \\ [\mathbf{V}_2, \mathbf{T}] &= qr(\mathbf{V}), \end{aligned}$$

(i.e., $\mathbf{V}_2\mathbf{T} = \mathbf{V}$ with \mathbf{V}_2 orthogonal and \mathbf{T} upper triangular matrices). Obtain additional basis vectors \mathbf{W}_2 via

$$\mathbf{B}\mathbf{V}_2 = \mathbf{W}_1\mathbf{R}_{12} + \mathbf{W}_2\mathbf{R}_{22} \quad \text{with } \mathbf{W}_1^H\mathbf{W}_2 = 0, \quad \mathbf{W}_2^H\mathbf{W}_2 = \mathbf{I}_k.$$

Then compute $\mathbf{H}_{12}, \mathbf{H}_{21}, \mathbf{H}_{22}, \mathbf{F}_1^+$ and \mathbf{F}_2 such that

$$\begin{aligned} \mathbf{A}\mathbf{V}_1 &= \mathbf{W}_1\mathbf{H}_{11} + \mathbf{W}_2\mathbf{H}_{21} + \mathbf{F}_1^+, \\ \mathbf{A}\mathbf{V}_2 &= \mathbf{W}_1\mathbf{H}_{12} + \mathbf{W}_2\mathbf{H}_{22} + \mathbf{F}_2. \end{aligned}$$

Finally, apply the QZ method (say) to the pair (\mathbf{H}, \mathbf{R}) to obtain unitary matrices \mathbf{Q}, \mathbf{Z} , an upper-triangular \mathbf{H}^+ and an upper triangular matrix \mathbf{R}^+ such that

$$\begin{aligned} \mathbf{H}\mathbf{Q} &= \mathbf{Z}\mathbf{H}^+, \\ \mathbf{R}\mathbf{Q} &= \mathbf{Z}\mathbf{R}^+, \end{aligned}$$

where $\mathbf{H} = (\mathbf{H}_{ij})$ and $\mathbf{R} = (\mathbf{R}_{ij})$, $i = 1, 2; j = 1, 2$, with the best approximations to the desired eigenvalues appearing as eigenvalues of $(1, 1)$ block of the pair $(\mathbf{H}^+, \mathbf{R}^+)$. Now, update

$$\begin{aligned} \mathbf{V}_1 &\leftarrow [\mathbf{V}_1, \mathbf{V}_2]\mathbf{Q}(:, 1:k), \quad \mathbf{W}_1 \leftarrow [\mathbf{W}_1, \mathbf{W}_2]\mathbf{Z}(:, 1:k), \\ \mathbf{H}_{11} &\leftarrow \mathbf{H}^+(1:k, 1:k), \quad \mathbf{R}_{11} \leftarrow \mathbf{R}^+(1:k, 1:k), \\ \mathbf{F}_1 &\leftarrow [\mathbf{F}_1^+, \mathbf{F}_2]\mathbf{Q}(:, 1:k). \end{aligned}$$

In this development, $p(\mathbf{A}, \mathbf{B})$ represents a matrix polynomial in \mathbf{A} and \mathbf{B} generated by a (preconditioned) iterative method designed to solve

$$(\mathbf{A} - \theta\mathbf{B})\hat{\mathbf{V}} = \mathbf{F}_1.$$

In fact, $\mathbf{G} \equiv p(\mathbf{A}, \mathbf{B})\mathbf{F}_1$ could easily represent a much more general object with each column of \mathbf{G} representing a separate iterative solution of the form

$$\mathbf{g}_j \approx (\mathbf{A} - \theta_j\mathbf{B})^{-1}\mathbf{F}_1\mathbf{y}_j, \quad j = 1, 2, \dots, k.$$

This could be made very efficient in terms of data movement per matrix-vector product. Each separate column would need two operations of the form $\mathbf{A}\mathbf{g}_j$ and $\mathbf{B}\mathbf{g}_j$. For example, a Richardson's iteration could take the form

```

G = F1 Y;
for j = 1, 2, ...
  G ← G $\Gamma$  - AG - BG $\Theta$ ;
end
  
```

where $\Gamma \equiv \text{diag}(\gamma_1, \gamma_2, \dots, \gamma_k)$ with reciprocal Richardson parameters γ_j and $\Theta \equiv \text{diag}(\theta_1, \theta_2, \dots, \theta_k)$ and $\mathbf{Y} \equiv [\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k]$ the current Ritz approximations to desired eigenvalues and vectors, i.e. $\mathbf{H}\mathbf{Y} = \mathbf{R}\mathbf{Y}\Theta$.

We may express the above discussion formally as the algorithm *BLKQZ* shown in Figure 6.2.

7. Computational Results and Conclusions. We shall present some very preliminary computational results to give some indication of the relative performance of three methods: *TFQZ*, *TBQZ*, *BLKQZ*. The purpose of these results is mainly to indicate that the methods have been programmed and will solve a difficult problem. There are many implementation details to consider and a number of parameter choices to be made. A thorough computational study including comparison with other methods is certainly called for.

Our results will consist of a comparison of the three methods on a single problem. The problem we consider is a symmetric generalized problem from the Harwell-Boeing collection. The matrices are stiffness and mass matrices were obtained through the Matrix Market from

<http://math.nist.gov/MatrixMarket/data/Harwell-Boeing/bcsstruc1/>

to form a generalized eigenvalue problem $\mathbf{A}\mathbf{x} = \mathbf{B}\mathbf{x}\lambda$. The matrix \mathbf{A} is BCSSTK12 and the matrix \mathbf{B} is BCSSTM12 from the BCSSTRUC1 set. BCSSTK12 and BCSSTM12 represent the consistent mass formulation for an ore car model. The consistent mass formulation leads to a non-diagonal mass matrix. All computations were done in Matlab Version 5.1.0.421 on a Sun SparcStation 20 Model 61 with 64 megabytes of RAM.

For these matrices, $n = 1473$ and \mathbf{A} has 17857 nonzero entries. The smallest four generalized eigenvalues are

```

3.469305448042201e+03
3.670875662014555e+03
5.538220410502827e+03
6.410197662646212e+03
  
```

and the largest generalized eigenvalue is on the order of 6.55e+08.

BLKQZ: Block Inexact QZ Process

Input: (A, B, V_1, k) such that $V_1^H V_1 = I_k$,

Output: (V_1, H_{11}, R_{11}) such that $AV_1 = W_1 H_{11}$, $V_1^T V_1 = I_k$.

$BV_1 = W_1 R_{11}$, $W_1^H W_1 = I_k$,

with H_{11} upper upper triangular and R_{11} upper triangular.

1. $\hat{W}_1 = BV_1$; $[W_1, R_{11}] = qr(\hat{W}_1)$;
 2. $F_1 \leftarrow AV_1$; $H_{11} \leftarrow (W_1^T F_1)$; $F_1 \leftarrow F_1 - W_1 H_{11}$;
 3. **for** $j = 1, 2, 3, \dots, k - 1$
 - 3.1. $\hat{V}_2 = \text{itsol}(A, B, M, F_1, Y_1)$;
 - 3.2. $S \leftarrow V_1^H \hat{V}_2$; $\hat{V}_2 \leftarrow \hat{V}_2 - V_1 S$;
 - 3.3. $[V_2, S] = qr(V_2)$;
 - 3.4. $\hat{W}_2 \leftarrow BV_2$; $R_{12} \leftarrow W_1^T W_2$;
 - 3.5. $\hat{W}_2 \leftarrow \hat{W}_2 - W_1 R_{12}$; $[W_2, R_{22}] = qr(\hat{W}_2)$;
 - 3.6. $H_{21} \leftarrow W_2^H F_1$; $F_1 \leftarrow F_1 - W_2 H_{21}$;
 - 3.7. $F_2 \leftarrow AV_2$; $H_{12} \leftarrow W_1^H F_2$;
 - 3.8. $F_2 \leftarrow F_2 - W_1 H_{12}$; $H_{22} = W_2^H F_2$; $F_2 \leftarrow F_2 - W_2 H_{22}$;
 - 3.9. $H \leftarrow \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix}$; $R \leftarrow \begin{pmatrix} R_{11} & R_{12} \\ 0 & R_{22} \end{pmatrix}$;
 - 3.10. $[Q, Z, H, R] = \text{qziter}(H, R, \text{'sort'})$;
 - 3.11. $V_1 \leftarrow [V_1, V_2]Q[:, 1:k]$; $W_1 \leftarrow [W_1, W_2]Z[:, 1:k]$;
 - 3.12. $F_1 \leftarrow [F_1, F_2]Q[:, 1:k]$;
 - 3.13. $H_{11} \leftarrow H(1:k, 1:k)$; $R_{11} \leftarrow R(1:k, 1:k)$;
4. **end**

FIG. 6.2. A Block Inexact QZ Process

Here, we list estimates of the computational and storage costs of the three routines and indicate the performance of each of them on this test problem. The term ‘‘matvec’’ stands for a matrix-vector product and the term ‘‘LU-solve’’ stands for solving the two successive triangular linear systems first with L and then with U as coefficient matrices.

TBQZ:

For a k -step factorization, the work and storage required for **TBQZ** is

- Storage: $2n(k + 1)$ plus storage for A, B, L, U
- Initial work:
 - 1 sparse LU-factorization,
 - $k + 1$ LU-solves,
 - $4n(k + 1)^2$ flops for orthogonalization.
- Work per iteration:
 - 1 LU-solve,
 - 1 matvec with (A, B) ,
 - $4n(k + 1)^2$ flops for orthogonalization,
 - sparse LU-factorization if there is a shift change.

For our run, $k = 9$ and the iteration was halted after four Ritz values had converged. The

TABLE 7.1
Eigenvalues calculated by *TBQZ*.

Eigenvalues	Error/ λ_{min}	Error/ λ_{max}
3.469305448324274e+03	2.8e-11	4.3e-16
3.670875661790737e+03	2.27e-11	3.4e-16
5.538220406841684e+03	3.7e-10	5.5e-15
6.410197672779293e+03	1.0e-09	1.5e-14

TABLE 7.2
Eigenvalues calculated by *TFQZ*.

Eigenvalues	Error/ λ_{min}	Error/ λ_{max}
3.469305447658971e+03	3.8e-11	5.8e-16
3.670875661610020e+03	4.1e-11	6.1e-16
5.538220410338460e+03	1.6e-11	2.5e-16
6.410197662356884e+03	2.9e-11	4.4e-16

code took 14 iterations and 7 matrix factorizations. The eigenvalues computed by *TBQZ* are shown in Table (7.1).

TFQZ:

For an m step factorization that retains a k step factorization after each implicit restart, the work and storage required is

- Storage: $2nm$ plus storage for **A**, **B**, **L**, **U**
- Initial work:
 - 1 sparse LU-factorization,
 - m LU-solves,
 - $4nm^2$ flops for orthogonalization.
- Work per iteration:
 - $m - k$ LU-solves,
 - $m - k$ matvecs with (A, B) ,
 - $4nm^2$ flops for orthogonalization,

For our run, $k = 4$ and $m = 12$ with $tol = 1.0e - 09$. The code took two iterations and 20 LU-solves. The eigenvalues computed by *TFQZ* are shown in Table (7.2).

BLKQZ:

The work and storage required with blocksize k is

- Storage: $4n(2k)$ plus storage for **A**, **B**, **L**, **U**
- Initial work:
 - 1 incomplete sparse LU-factorization,
 - 1 block ILU-solve,
 - $4n(2k)^2$ flops.
- Work per iteration:
 - 1 block ILU-solve,
 - 1 block matvec with (A, B) ,
 - $30n(2k)^2$ flops,

TABLE 7.3
Eigenvalues calculated by BLKQZ.

Eigenvalues	Error/ λ_{min}	Error/ λ_{max}
3.469305447907588e+03	1.4e-11	2.0e-16
3.670875661903084e+03	1.1e-11	1.7e-16
5.538220410459639e+03	4.4e-12	6.6e-17
6.410197662585929e+03	6.1e-12	9.2e-17

For our run, $k = 4$. The code took 43 matrix accesses, 43 block matvecs (A,B) and 443 individual matrix-vector products. The eigenvalues computed by *BLKQZ* are shown in Table (7.3).

In each routine, we used a reference shift of $\sigma = 3.4e+3$ and in the call to `tfqz` we passed $A - \sigma B$ in place of B and B in place of A in the calling sequence. This is mathematically equivalent to using implicit restarting with the shift-invert operator $(A - B)^{-1}B$ and the convergence results confirm that. For the *BLKQZ* method we used a block variant of BICGSTAB that we constructed from the single vector code in the templates collection [1] and with an incomplete LU preconditioner from Matlab. We were able to arrange the code so that each column of the right hand side represented a residual of the form

$$\mathbf{r}_j = (A - \mu_j B)\mathbf{x}_j$$

but used the same preconditioner for the whole block. Typically, not all of the column equations converged and our cut off was 10 iterations. As the results show, this was sufficient for convergence.

With these results, it is difficult to choose between the methods. Here, *TFQZ* seems to be the winner but that is in absence of any architecture considerations and without specific comparison between ILU and complete LU costs. We did not report flop counts or timings because the implementations are fairly crude at this point in time. These results only indicate that the three methods are indeed implementable and that they work on a challenging problem.

The real value of the *TBQZ* may lie in its applicability to rational interpolation with respect to constructing reduced order models of state space control systems as explored in [9]. More investigation and testing needs to be done with respect to shift selection and selecting the right hand side of the BQZ equations. The preconditioned *BLKQZ* is very promising with respect to parallel performance but is far from robust at this time.

Acknowledgement. The author would like to thank R. B. Lehoucq and C. Yang for several discussions and comments. He would also like to thank the anonymous referee for several useful comments and for catching some mistakes in the section on inexact iterations.

REFERENCES

[1] RICHARD BARRETT, MICHAEL BERRY, TONY F. CHAN, JAMES DEMMEL, JUNE DONATO, JACK DONGARRA, VICTOR EIJKHOUT, ROLDAN POZO, CHARLES ROMINE, AND HENK VAN DER VORST, *Templates for the Solution of Linear Systems: Building Blocks for Iterative Methods*, SIAM, Philadelphia, USA, 1993.

[2] M. CROUZEIX, B. PHILIPPE AND M. SADKANE, *The Davidson method*, SIAM J. Sci. Comput., 15 (1994), pp. 62–76.

[3] J. DANIEL, W. B. GRAGG, L. KAUFMAN AND G. W. STEWART, *Reorthogonalization and stable algorithms for updating the Gram–Schmidt QR factorization*, Math. Comp., 30 (1976), pp. 772–795.

- [4] ERNEST R. DAVIDSON, *The iterative calculation of a few of the lowest eigenvalues and corresponding eigenvectors of large real-symmetric matrices*, J. Comput. Phys., 17 (1975), p. 87.
- [5] T. ERICSSON AND A. RUHE, *The spectral transformation Lanczos method for the numerical solution of large sparse generalized symmetric eigenvalue problems*, Math. Comp., 35 (1980), pp. 1251–1268.
- [6] D.R. FOKKEMA, G.L.G. SLEIJPEN AND H.A. VAN DER VORST, *Jacobi-Davidson style QR and QZ algorithms for the partial reduction of matrix pencils*, Technical Report Preprint 941, Department of Mathematics, Utrecht University, Utrecht, The Netherlands, January 1996.
- [7] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, Johns Hopkins University Press, 3rd Ed., Baltimore, 1996.
- [8] R. G. GRIMES, J. G. LEWIS AND H. D. SIMON, *A shifted block Lanczos algorithm for solving sparse symmetric generalized eigenproblems*, SIAM J. Matrix Anal. Appl., 15(1) (1994), pp. 228–272.
- [9] E.J. GRIMME, *Krylov Projection Methods for Model Reduction*, Ph.D. thesis, University of Illinois, Urbana-Champaign, 1997.
- [10] A.V. KNYAZEVR, *Convergence rate estimates for iterative methods for a mesh symmetric eigenvalue problem*, Sov. J. Numer. Anal. Math. Modeling, 2(5) (1987), pp. 371–396.
- [11] R. B. LEHOUCQ AND KARL MEERBERGEN, *Using generalized Cayley transformations within an inexact rational Krylov sequence method*, SIAM J. Matrix Anal. Appl., (1998), to appear.
- [12] K. MEERBERGEN AND A. SPENCE, *Implicitly restarted Arnoldi with purification for the shift–invert transformation*, Math. Comp., 218 (1997), pp. 667–689.
- [13] C.B. MOLER AND G. W. STEWART, *An algorithm for generalized matrix eigenvalue problems*, SIAM J. Numer. Anal., 10 (1973), pp. 241–256.
- [14] R. B. MORGAN AND D. S. SCOTT, *Generalizations of Davidson’s method for computing eigenvalues of sparse symmetric matrices*, SIAM J. Sci. Statist. Comput., 7 (1986), pp. 817–825.
- [15] J. OLSEN, P. JORGENSEN AND J. SIMONS, *Passing the one-billion limit in full configuration interaction (fci) calculations*, Chem. Phys. Letters, 169(6) (1990), pp. 463–472.
- [16] A. RUHE, *Rational Krylov sequence methods for eigenvalue computations*, Linear Algebra Appl., 58 (1984), pp. 391–405.
- [17] G. DE SAMBLANX, K. MEERBERGEN AND A. BULTHEEL, *The implicit application of a rational filter in the RKS method*, BIT, 37 (1997), pp. 925–947.
- [18] G. L. G. SLEIJPEN AND H.A. VAN DER VORST, *A Jacobi-Davidson iteration method for linear eigenvalue problems*, SIAM J. Matrix Anal. Appl., 17(2) (1996), pp. 401–425.
- [19] G.L.G. SLEIJPEN, J.G.L. BOOTEN, D.R. FOKKEMA AND H.A. VAN DER VORST, *Jacobi-Davidson type methods for generalized eigenproblems and polynomial eigenproblems*, BIT, 36(3) (1996), pp. 595–633.
- [20] D. C. SORENSEN, *Implicit application of polynomial filters in a k-step Arnoldi method*, SIAM J. Matrix Anal. Appl., 13(1) (1992), pp. 357–385.
- [21] D.C. SORENSEN AND C. YANG, *A truncated RQ-iteration for large scale eigenvalue calculations*, SIAM J. Matrix Anal. Appl., to appear.